



HEADS UP CONCUSSION IN YOUTH SPORTS

A Fact Sheet for COACHES

To download the coaches fact sheet in Spanish, please visit www.cdc.gov/ConcussionInYouthSports
Para descargar la hoja informativa para los entrenadores en español, por favor visite:
www.cdc.gov/ConcussionInYouthSports

THE FACTS

- A concussion is a **brain injury**.
- All concussions are **serious**.
- Concussions can occur **without** loss of consciousness.
- Concussions can occur **in any sport**.
- Recognition and proper management of concussions when they **first occur** can help prevent further injury or even death.

WHAT IS A CONCUSSION?

A concussion is an injury that changes how the cells in the brain normally work. A concussion is caused by a blow to the head or body that causes the brain to move rapidly inside the skull. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious. Concussions can also result from a fall or from players colliding with each other or with obstacles, such as a goalpost.

The potential for concussions is greatest in athletic environments where collisions are common.¹ Concussions can occur, however, in **any** organized or unorganized sport or

recreational activity. As many as 3.8 million sports- and recreation-related concussions occur in the United States each year.²

RECOGNIZING A POSSIBLE CONCUSSION

To help recognize a concussion, you should watch for the following two things among your athletes:

1. A forceful blow to the head or body that results in rapid movement of the head.

-and-

2. Any change in the athlete’s behavior, thinking, or physical functioning. (See the signs and symptoms of concussion listed on the next page.)

It's better to miss one game than the whole season.

ACTION PLAN

WHAT SHOULD A COACH DO WHEN A CONCUSSION IS SUSPECTED?

- 1. Remove the athlete from play.** Look for the signs and symptoms of a concussion if your athlete has experienced a bump or blow to the head. Athletes who experience signs or symptoms of concussion should not be allowed to return to play. When in doubt, keep the athlete out of play.
- 2. Ensure that the athlete is evaluated right away by an appropriate health care professional.** Do not try to judge the severity of the injury yourself. Health care professionals have a number of methods that they can use to assess the severity of concussions. As a coach, recording the following information can help health care professionals in assessing the athlete after the injury:
 - Cause of the injury and force of the hit or blow to the head
 - Any loss of consciousness (passed out/knocked out) and if so, for how long
 - Any memory loss immediately following the injury
 - Any seizures immediately following the injury
 - Number of previous concussions (if any)
- 3. Inform the athlete's parents or guardians about the possible concussion and give them the fact sheet on concussion.** Make sure they know that the athlete should be seen by a health care professional experienced in evaluating for concussion.
- 4. Allow the athlete to return to play only with permission from a health care professional with experience in evaluating for concussion.** A repeat concussion that occurs before the brain recovers from the first can slow recovery or increase the likelihood of having long-term problems. Prevent common long-term problems and the rare *second impact syndrome* by delaying the athlete's return to the activity until the player receives appropriate medical evaluation and approval for return to play.

REFERENCES

1. Powell JW. Cerebral concussion: causes, effects, and risks in sports. *Journal of Athletic Training* 2001; 36(3):307-311.
2. Langlois JA, Rutland-Brown W, Wald M. The epidemiology and impact of traumatic brain injury: a brief overview. *Journal of Head Trauma Rehabilitation* 2006; 21(5):375-378.
3. Lovell MR, Collins MW, Iverson GL, Johnston KM, Bradley JP. Grade 1 or "ding" concussions in high school athletes. *The American Journal of Sports Medicine* 2004; 32(1):47-54.
4. Institute of Medicine (US). Is soccer bad for children's heads? Summary of the IOM Workshop on Neuropsychological Consequences of Head Impact in Youth Soccer. Washington (DC): National Academy Press; 2002.
5. Centers for Disease Control and Prevention (CDC). Sports-related recurrent brain injuries-United States. *Morbidity and Mortality Weekly Report* 1997; 46(10):224-227. Available at: www.cdc.gov/mmwr/preview/mmwrhtml/00046702.htm.

If you think your athlete has sustained a concussion... take him/her out of play, and seek the advice of a health care professional experienced in evaluating for concussion.